



# Effect of Various Silicon Sources on Nutrient Uptake in Rice

Guntamukkala Babu Rao<sup>1\*</sup>, Poornima Yadav P<sup>2</sup> and Elizabeth K Syriac<sup>3</sup>

Department of Agronomy  
Kerala Agricultural University, Thiruvananthapuram 680654 (Kerala)

## ABSTRACT

Silicon nutrition is gaining importance in agriculture owing to its positive effects in rice production. A field experiment was carried out at the farmer's field in Kerala during Kharif, 2016 to evaluate different silicon sources on nutrient uptake by rice and available nutrient status of soil after the harvest. The experiment was laid out in randomised block design with seven treatments and three replications. Different silicon sources viz., potassium silicate, fine silica, rock dust, rice husk ash were involved in the treatments and fertilizer application was done according to the recommended dose of fertilizers as per Package of Practice of Kerala Agricultural University. Silicon nutrition have shown significant influence on the total nutrient uptake by the crop and available soil nutrient status of soil after the harvest. Among the treatments, the Treatment T<sub>6</sub> i.e., fine silica @ 50 kg/ha + rice husk ash @ 250 kg/ha, has shown the better results with respect to nutrient uptake {N (189.74 kg/ha); P (17.19 kg/ha); K (127.38 kg/ha) and Si (345.14 kg/ha)} by the crop and available nutrient status of the soil {N (377.73 kg/ha); P (36.37 kg/ha); K (206.25 kg/ha) and Si (83.61 kg/ha)} after the harvest.

**Key Words:** Laterite soils, Nutrient Uptake, Nutrient status, Rice, Silicon.

## INTRODUCTION

Rice is the main staple food crop of Kerala. The rice farming sector of Kerala is facing a series of problems which leads to a drastic decline in production due to numerous soil related constraints like Iron and Aluminium toxicity and high acidity of the soils of the state (Maneesha et al, 2016). Majority of Kerala soils are lateritic in nature which are low in Organic carbon, N and K, very low in Ca and Mg coupled with the iron and aluminium toxicities which need separate management package (Anonymous, 2016). The foremost reasons behind the low productivity of rice in laterite soils of Kerala mainly in lowland situations are due to the low nutrient status of the soil

Silicon (Si) is one of the abundant element in the earth's crust which is known to have several beneficial effects on crop growth, especially for Poaceae crops like Rice (Devanur, 2015; Rao, 2017). The potential of silicon in enhancing rice yield has been demonstrated in numerous studies especially under biotic and abiotic stress conditions

viz. diseases, insect pests, drought, salinity, heavy metals (Epstein, 2001). Silicon is known to reduce the concentration of toxic elements like Fe, Al, other heavy metals in laterite derived paddy soils and also improve soil physical properties and available nutrient status in soil (Devanur, 2015). Rice being a high Si demanding crop, increasing rice yield per unit area is relatively associated with Silicon depletion, which is a matter of concern. In general farmers export silicon from field by removing straw residues with the harvest and the exogenous application of silicon in rice is overlooked. Therefore, a continued supply of silicon would be required predominantly for the healthy and productive development of plant during all growth stages (Epstein, 2001). Hence, the present investigation was undertaken with an objective to assess the effect of silicon nutrition in rice on total nutrient uptake by the crop and available nutrient status of soil after the harvest.

## MATERIALS AND METHODS

The field experiment was conducted at a farmer's field in Kerala, during Kharif 2016. The

\*Corresponding Author's Email: sekhar2486@gmail.com

<sup>1</sup>M.Sc. Scholar. <sup>2</sup>Assistant Professor. Krishi Vigyan Kendra, Kottarakkara, Kollam, Kerala and <sup>3</sup>Professor (Agronomy).

soil of the experimental site was sandy clay loam, acidic in nature (pH 4.50), high in OC (1.01%) and safe EC (0.10 dS/m). The initial nutrient status of the soil were N (550.5 kg/ha), P (16.86 kg/ha), K (196.90 kg/ha) and Si (45.02 kg/ha) respectively. The experiment was laid out in randomized block design with seven treatments and three replications with each plot size of 5 m x 4 m using Rice variety Uma, which was transplanted during first week of July with a spacing of 20 x 15 cm. All treatments were supplied with similar recommended dose of fertilizers i.e., Lime @ 150 kg/ha + farm yard manure @ 5 t/ha + NPK @ 90:45:120 kg/ha (KAU, 2016) The treatments were, T<sub>1</sub>-Fine silica @ 100 kg/ha; T<sub>2</sub>-Fine silica @ 75 kg/ha + rock dust @ 25 kg/ha; T<sub>3</sub>-Fine silica @ 75 kg/ha + foliar application of K<sub>2</sub>SiO<sub>3</sub> at maximum tillering stage @ 0.5%; T<sub>4</sub>-Fine silica @ 50 kg/ha + rock dust @ 25 kg/ha + foliar application of K<sub>2</sub>SiO<sub>3</sub> at maximum tillering stage @ 0.5%; T<sub>5</sub>-Fine silica @ 75 kg/ha + rice husk ash @ 125 kg/ha; T<sub>6</sub>-Fine silica @ 50 kg/ha + rice husk ash @ 250 kg/ha; T<sub>7</sub>- Fine silica @ 50 kg/ha + rice husk ash @ 125 kg/ha + foliar application of potassium silicate at maximum tillering stage @ 0.5%. Silicon sources such as fine silica, rock dust, and rice husk ash were applied basally as per treatments at transplanting, and foliar application of potassium silicate @ 0.5 % at maximum tillering stage. Soil samples were analysed for available nutrient status before and after the harvest of the crop and expressed as kg/ha. Plant samples were collected at harvest stage and analyzed for different nutrients viz., N, P, K and Si. The total uptake of N, P, K and Si by the plant at harvest was calculated as the product of the respective nutrient content and plant dry weight and expressed as kg/ha. The data obtained were subjected to statistical analysis and were tested at five per cent level of significance to interpret the treatment differences.

## RESULTS AND DISCUSSION

### Effect of silicon nutrition on nutrient content in rice straw, grain and total nutrient uptake

The silicon nutrition had no shown significant

influence on N content in grain and straw, but total N uptake increased significantly by silicon nutrition. With respect to total N uptake, T<sub>6</sub> (Fine silica @ 50 kg/ha + rice husk ash @ 250 kg/ha) was superior with an uptake of 189.74 kg/ha. The available N content of soil was also low for the above treatment after harvest compared to the initial soil N. This might naturally be due to enhanced absorption of N by the crop ultimately leading to higher N uptake by plant, resulting in low available N status in soils. Similar results have also been reported by Rao (2018) and Devanur (2015).

Phosphorus concentration in plant and uptake of P were positively influenced by silicon application. Treatment T<sub>6</sub> (fine silica @ 50 kg/ha + rice husk ash @ 250 kg/ha) produced significantly higher content of P in grain and total P uptake, this treatment received the highest quantity of silicon i.e. 200 kg/ha. The P content of straw was also found to be the highest in T<sub>6</sub> (0.07 %), followed by T<sub>7</sub> (0.06 %). The available P content in the soil after the experiment was also high in the above treatments. The monosilicic acid anions released from silicon sources might have replaced the phosphate anions released from Fe and Al phosphate, which might have resulted in higher P content and uptake by the plants. Increase in P uptake by the rice crop increased from 26 to 34 per cent when P as single superphosphate was applied along with a silicate fertilizer. Tavakkoli *et al.* (2011) reported that overall beneficial effect of silicon may be attributed to a higher P: Mn ratio in the plant shoot due to the decreased Mn and Fe uptake, and thus indirectly improving P utilization within the rice plants. Addition of silicon fertilizers also increased the pH in acid soils which will release P from Fe-P and Al-P complexes (Rao, 2017)

The data with respect to K content in grain, straw and total K uptake are presented in Table 2. The content of K in grain, straw and total uptake of K by rice crop increased with silicon application. Fine silica @ 50 kg/ha + rice husk ash @ 125 kg/ha + foliar application of potassium silicate at maximum tillering stage @ 0.5 % spray (T<sub>7</sub>) and

## Effect of Various Silicon Sources

**Table1. Effect of silicon nutrition on the N and P content in grain, straw and total N and P uptake by Rice.**

Treatment	N content (%)		Total N uptake (kg/ha)	P content (%)		Total P uptake (kg/ha)
	Grain	Straw		Grain	Straw	
T <sub>1</sub>	1.46	0.97	166.73	0.15	0.04	12.08
T <sub>2</sub>	1.45	1.01	164.49	0.14	0.03	10.39
T <sub>3</sub>	1.45	0.95	166.02	0.14	0.04	11.52
T <sub>4</sub>	1.42	1.03	169.30	0.15	0.05	12.86
T <sub>5</sub>	1.51	0.97	175.25	0.16	0.06	14.49
T <sub>6</sub>	1.55	1.04	189.74	0.18	0.07	17.19
T <sub>7</sub>	1.56	0.99	181.36	0.17	0.06	15.44
S E m±	0.096	0.051	4.582	0.000	0.000	0.112
CD (0.05)	NS	NS	9.985	0.005	0.009	0.243

fine silica @ 50 kg/ha + rice husk ash @ 250 kg/ha (T<sub>6</sub>) were significantly superior with respect to K content in straw. However, T<sub>7</sub> was found to be the treatment with highest grain K and total K uptake. Soil application of silicon has synergistic interaction with applied K and also promotes the release of K from the exchange sites to the soil solution by the hydrogen ions produced during the oxidation of Fe and Al compounds. Silicon application increased yield response to applied potassium in upland rice. Similar beneficial effect of silicon fertilizers on K content in plant and K uptake are reported by Singh and Singh (2005) and Sunil kumar (2000).

### Effect of silicon nutrition on Silicon content in rice straw, grain and total uptake

The data with respect to Si content in grain, straw and total Si uptake are presented in Table 2. The silicon nutrition of rice evaluated in terms of concentration and uptake of silicon was influenced by silicon fertilization. With respect to silicon content in grain and straw, T<sub>7</sub> and T<sub>6</sub> were significantly superior to other treatments. Silicon supply in T<sub>7</sub> was less, but foliar application of potassium silicate helped to improve silicon uptake. However, with respect to available silicon in soil, T<sub>6</sub> was superior

to T<sub>7</sub>. The increase in plant available silicon in the soil was usually accompanied by increased silicon accumulation in the plant, which might have result in increased growth and productivity in several crops, especially rice. Silicon content of rice straw shows large variations from 1.7 to 9.3% and is influenced by several factors such as soil, irrigation water quality, amount of fertilizers applied, rice cultivars and season. The straw silica content of rice at harvest ranged from 4.8 to 13.5% in dry season and from 4.3 to 10.3%, in wet season (Devanur, 2015).

### Effect of silicon nutrition on available nutrient status of soil

The data on available nutrient (N, P, K and Si) content in soil are presented in Table 3. The available N content in the soil was not significantly influenced by the treatments. The treatments had not shown significant effect on available N in soil, but when compared to initial soil N status, there was a decline in soil N status in all the treatments. This decrease in available N in soil might be due to enhanced uptake of soil N, because silicon in soil has the ability to raise the optimum N rate, thus enhancing the productivity of existing lowland paddy fields (Rao, 2017).

**Table 2. Effect of silicon nutrition on the K and Si content in grain, straw and total K and Si uptake by Rice.**

Treatment	K content (%)		Total K uptake (kg/ha)	Si content (%)		Total Si uptake (kg/ha)
	Grain	Straw		Grain	Straw	
T <sub>1</sub>	0.40	0.95	105.81	0.76	3.20	322.78
T <sub>2</sub>	0.36	0.93	99.82	0.61	2.8	273.89
T <sub>3</sub>	0.39	0.97	107.25	0.68	3.11	311.19
T <sub>4</sub>	0.37	0.95	104.84	0.59	2.9	282.27
T <sub>5</sub>	0.50	0.96	113.10	0.85	3.21	330.11
T <sub>6</sub>	0.56	1.06	127.38	0.92	3.29	345.14
T <sub>7</sub>	0.86	1.09	147.07	0.96	3.37	352.93
S E m±	0.115	0.036	1.414	0.051	0.057	2.192
CD (0.05)	0.059	0.082	3.082	0.108	0.125	4.778

The available P content in soil was significantly higher in the T<sub>6</sub>, followed by T<sub>7</sub> and T<sub>5</sub>. This increase in P might be due to the possibility of replacing the phosphate anion [HPO<sub>4</sub>]<sup>2-</sup> from Al and Fe phosphates by monosilicic acid [Si(OH)<sub>3</sub>]<sup>-</sup> of silicon sources. Guntzer *et al.* (2012) observed that there was an increase in the response of applied phosphorus in rice, when applied along with silicon fertilizers.

The available potassium content in soil was significantly influenced by the silicon application. The highest available K was found in T<sub>6</sub>, which was followed by T<sub>7</sub>, T<sub>1</sub>, T<sub>5</sub> and T<sub>4</sub>. The production of hydrogen ions during reduction of Fe and Al might have helped in the release of K from the exchange sites or from the fixed pool to the soil solution. Yadav (2017) stated that beside yield enhancement in rice, silicon also has many fold advantages of increasing availability of major nutrients and also alleviating iron toxicity problems in soils. These results were confirmative with the findings of and Mali and Aery (2008).

Silicon nutrition significantly influenced soil silicon status also. The soil silicon was found to be higher in all the treatments after harvest compared to the initial status, but the highest soil available

silicon was found in T<sub>6</sub>, followed by T<sub>5</sub>. The silicon applied through various silicon sources, would have prevailed in soil as monosilicic acid (H<sub>4</sub>SiO<sub>4</sub>) due to its residual activity and enhanced soil silicon availability. These findings were in agreement with those reported by Singh *et al* (2006) and Korndorfer *et al* (2001). Prasanta and Heinz (2009) reported that changes in the pH of soils due to soil flooding significantly influence the solubility of Fe, P and Si in soil; so also plant available soil silicon increases due to increase in soil reaction. In the present study also, the increase in soil reaction compared to the initial value might have resulted in significantly higher silicon content in soil.

## CONCLUSION

The silicon in soil is decreasing at a rapid rate due to intensive cultivation of Rice and also due to lack of incorporation of rice crop residues into the soil. In general, the farmers remove all the rice stubbles after the harvest of the crop in order to prepare the field for next crop. By this process soil silicon is not getting replenished. Nowadays the importance of silicon is much felt by researches as it have many advantages in the rice crop production. To solve the deficiency of silicon in soil, the rice stubbles should be incorporated and along with that

## Effect of Various Silicon Sources

**Table 3. Effect of silicon nutrition on available nutrients (N, P, K and Si) in soil.**

Treatment	Available nutrients (kg/ha)			
	N	P	K	Si
T <sub>1</sub>	356.10	32.58	199.10	75.40
T <sub>2</sub>	323.33	31.87	183.40	74.92
T <sub>3</sub>	363.43	29.54	177.87	74.17
T <sub>4</sub>	315.53	27.49	187.84	74.06
T <sub>5</sub>	370.99	34.72	196.06	80.76
T <sub>6</sub>	377.73	36.37	206.25	83.61
T <sub>7</sub>	366.68	33.96	204.89	79.33
S E m±	19.971	1.567	8.675	1.547
CD (0.05)	NS	3.415	18.904	3.372

various organic and inorganic sources can be used. In the present research the treatments were sources like rice husk ash and fine silica were applied in-addition to the Package of practices followed by Kerala Agricultural University, resulted in the better nutrient uptake by rice and also the available nutrient status was also increased.

### REFERENCES

- Anonymous (2016). Soils of Kerala [On-line]. Available: <http://www.keralaagriculture.gov.in> [25 DEC.2016].
- Devanur V (2015). SILICON-Solution for tomorrow, Concept note. Available: <http://www.privilifesciences.com/download/silicon-supplement.pdf>. [25 Dec. 2016].
- Epstein E (2001). Silicon in plants. *Stud Plant Sci* 8: 1-15.
- Guntzer F, Keller C and Meunier J D (2012). Benefits of plant silicon for crops: A review. *Agron, Sustain Dev* 32: 201–213.
- KAU (Kerala Agricultural University). 2016. *Package of Practices Recommendations: Crops* (15<sup>th</sup> Ed.). Kerala Agricultural University, Vellanikkara, Thrissur, 393p.
- Korndorfer G H, Snyder G H, Ulloa M, Powell G and Datnoff L E (2001). Calibration of soil and plant silicon analysis for rice production. *J Plant Nutr* 24: 1071-1084.
- Mali M and Aery N C (2008). Silicon effects on nodule growth, dry matter production, and mineral nutrition of cowpea (*Vigna unguiculata*). *J Plant Nutr Soil Sci* 171: 835–40.
- Maneesh P and Deepa N R (2016). Trend analysis of area, production and productivity of rice in Kerala in the context of food security. *Int J Agric Res Rev* 4(8): 538-546.
- Prasanta K P and Heinz UN (2009). Dynamics of water soluble silica and silicon nutrition of rice in relation to changes in iron and phosphorus in soil solution due to soil drying and reflooding. *Archives Agron Soil Sci* 56(6): 605–622.
- Rao G B, Yadav P I and Syriac E K (2017). Silicon nutrition in rice- A review. *J. Pharmacognosy Phytochem* 6(6): 390-392.
- Rao G B, Yadav P I and Syriac E K (2018). Effect of silicon on soil physico-chemical properties in laterite derived paddy soils of Kerala. *J Krishi Vigyan* 6(2): 75-77.
- Singh K K and Singh K (2005). Effect of N and Si on growth, yield attributes and yield of rice in Alfisols. *Int Rice Res Notes* 12: 40–41.
- Singh K, Singh R, Singh J P, Singh Y and Singh K K (2006). Effect of level and time of silicon application on growth, yield and its uptake by rice. *Indian J Agric Sci* 76(7): 410-413.
- Sunilkumar B (2000). *Suitability of upland rice (Oryza sativa L.) cultivars for shaded situations*. M.Sc. (Ag) thesis, Kerala Agricultural University, Thrissur, 117p.
- Tavakkoli E H, English P and Guppy N C (2011). Silicon and phosphorus to mitigate manganese toxicity in rice in a highly weathered soil. *Commun Soil Sci Plant Analysis* 42: 503-513
- Yadav P. P I, Manu C R and Noble Abraham (2017). Silicon nutrition for sustainable rice production in iron toxic laterite soils of Kollam district in Kerala. *J. Krishi Vigyan* 5(2): 150-153.

Received on 08/06/2019

Accepted on 08-09-2019