



Impact of Rootstock Selection and Cane Regulation on the Organoleptic Characteristics of Wine

Suraj S Hanni^{1*}, S N Patil², Anand G . Nanjapanavar³, I B Biradar⁴, Mallikarjun Awati⁵, Venkateshalu⁶ and Pooja Murthy S⁷

Main Horticultural Research and Extension Centre
University of Horticultural Sciences, Bagalkot, 587104, (Karnataka), India

ABSTRACT

An investigation was conducted to assess the impact of rootstock selection and cane regulation on the organoleptic characteristics of wine in the semi-arid region of Karnataka. The study was carried out during 2022-2024 at the Main Horticultural Research and Extension Centre, Bagalkot, using a split-split plot design with two rootstocks (R₁: Dogridge and R₂: 110 R), five grape varieties (V₁: Cabernet Sauvignon, V₂: Medica, V₃: Shiraz, V₄: Sauvignon Blanc and V₅: Chenin Blanc), and three cane regulation levels (C₁: control, C₂: 36 canes/vine and C₃: 24 canes/vine). Organoleptic properties, including appearance, aroma, taste, and overall acceptability, were evaluated using a 20-point hedonic scale over two years and pooled data. Results revealed significant differences among treatments. Rootstock R₂ consistently recorded higher scores across all organoleptic parameters, with pooled appearance and aroma scores of 17.90 and 18.16, respectively, compared to R₁. Among varieties, V₂ exhibited superior performance, with the highest pooled appearance (17.47) and aroma (17.31) scores, followed by V₃ and V₁. Cane regulation influenced appearance but had a non-significant effect on aroma, with C₃ recording the highest pooled scores for appearance (15.78) and aroma (16.13). Interaction effects were significant, with the combination R₂V₂C₃ achieving the highest scores for both appearance (20.68) and aroma (20.05). The findings demonstrate that rootstock 110 R, Medica variety, and a 24-cane regulation approach significantly enhance the sensory quality of wine in semi-arid conditions. These results emphasize the importance of integrating rootstock selection and cane regulation to optimize wine quality. The study provided valuable insights for viticultural practices in similar agro-climatic regions, contributing to sustainable and high-quality wine production systems.

Keywords: Cane regulation, Organoleptic characteristics, Rootstock, Selection, Wine quality.

INTRODUCTION

Grape cultivation is considered to be an economic venture for the farmers due to its high monetary returns both from its fruits and processed products like wine (Kaur *et al*, 2021). The production of wine from fruits of grape is an intricate process, where the quality and sensory attributes of the final product are greatly influenced by viticultural practices. Organoleptic evaluation, encompassing the sensory parameters of appearance, aroma, taste, and mouthfeel, plays a crucial role in determining the acceptability and market value of wine. Factors such as rootstock selection and cane regulation are pivotal in shaping these sensory characteristics, as they influence the

growth, yield, and chemical composition of grape berries, which, in turn, impact the wine's quality and sensory profile. Rootstocks contribute to the vine's tolerance to biotic and abiotic stresses, nutrient uptake, and overall vine vigor (Singh *et al*, 2025), which directly affect the accumulation of phenolic compounds, organic acids, and sugars in the grape berries which are the key determinants of wine's organoleptic properties. Cane regulation, primarily through pruning intensity, alters the vine's canopy structure, light interception, and the balance between vegetative and reproductive growth. This influences the development of flavour precursors and secondary metabolites that enhance wine aroma, taste, and mouthfeel. (Kiran *et al*, 2019). This study was

Corresponding Author's Email - srj.hanni@gmail.com

1,7University of Horticultural Sciences, Bagalkot, 587104, (Karnataka), India

2,3Department of Fruit Science, University of Horticultural Sciences, Bagalkot, 587104, (Karnataka), India

4,5Department of N.R.M., University of Horticultural Sciences, Bagalkot, 587104, (Karnataka), India

6College of Horticulture Kolar, University of Horticultural Sciences, Bagalkot, 587104, (Karnataka), India

undertaken to assess the organoleptic properties of wine produced from five grape varieties grown on two rootstocks and subjected to three levels of cane regulation. By evaluating the appearance, aroma, taste, and overall acceptability of wine over two years and pooled data, the study aimed to identify significant differences and interactions among the treatments and determine the most favourable combinations for enhancing wine quality. The research holds immense relevance for viticulture practices in regions like Karnataka's northern dry zone, where managing environmental and climatic challenges is critical for producing high-quality wine. The findings are expected to provide valuable insights into optimizing rootstock selection and cane regulation to improve the sensory attributes of wine, contributing to the advancement of sustainable and high-quality wine production systems.

MATERIALS AND METHODS

The investigation on “Influence of rootstock and cane regulation on growth, yield, and quality of wine grapes” was conducted during 2022-2024 at MHREC, Block J1, UHS, Bagalkot (16° 9' N, 75° 37' E, 542 m MSL) in the northern dry zone of Karnataka. The experiment used a split-split plot design with 30 treatments replicated three times. Spacing was 9 ft × 4 ft. The main treatments were rootstocks (Dogridge and 110 R), sub-treatments were cultivars (Cabernet Sauvignon, Medica, Shiraz, Sauvignon Blanc and Chenin Blanc), and sub-sub treatments were cane regulation levels (C₁: Control, C₂: 36 canes/vine and C₃: 24 canes/vine). Data were recorded on randomly selected plants. Wine samples were organoleptically analyzed using a 20-point hedonic scale (appearance, aroma, taste, colour, overall acceptability) by 10 panel members, following standard wine-tasting procedures (Veena *et al*, 2015).

Hedonic scale:

18-20 excellent	15-17 good	12-14 fair		
9-11 ordinary	6-8 poor	3-5 bad	1-2 very bad	

RESULTS AND DISCUSSION

Organoleptic evaluation of appearance

The organoleptic evaluation of appearance in 2023, 2024, and in pooled data showed significant differences due to rootstocks (Table 1). In pooled data, rootstock R₂ recorded significantly higher appearance scores (17.90) compared to rootstock R₁ (12.94). The variation in appearance scores due to varieties was significant as well. Variety V₂ exhibited the highest

appearance score (17.47), followed by V₃ (16.46) and V₁ (15.55). The lowest appearance score was recorded by variety V₅ (12.91). Cane regulation also had a significant effect on appearance. In pooled data, cane C₃ showed the highest appearance score (15.78), followed by C₂ (15.40), and the control C₁ (15.09). The interaction between rootstocks, varieties, and cane regulation influenced the organoleptic evaluation of appearance significantly. In pooled data, the treatment combination R₂V₂C₃ recorded the highest appearance score (20.68), followed by R₂V₂C₂ (20.15) and R₂V₃C₃ (19.23).

Organoleptic evaluation of aroma

The aroma evaluation in 2023, 2024, and in pooled data showed significant differences due to rootstocks (Table 1). In pooled data, rootstock R₂ recorded significantly higher aroma scores (18.16) compared to rootstock R₁ (13.42). The variation in aroma scores due to varieties was significant. Variety V₂ had the highest aroma score (17.31), followed by V₃ (16.63) and V₁ (16.05). The lowest aroma score was recorded by variety V₅ (13.62). Cane regulation had a non-significant effect on aroma during 2023, 2024, and in pooled data. In pooled data, cane C₃ exhibited the highest aroma score (16.13), followed by C₂ (15.79) and control C₁ (15.44). The interaction between rootstocks, varieties, and cane regulation significantly influenced the aroma evaluation. In pooled data, the treatment combination R₂V₂C₃ recorded the highest aroma score (20.05), followed by R₂V₂C₂ (19.73) and R₂V₃C₃ (19.15).

Organoleptic evaluation of taste

The taste evaluation in 2023, 2024, and in pooled data revealed significant differences due to rootstocks (Table 2). In pooled data, rootstock R₂ recorded significantly higher taste scores (19.54) compared to rootstock R₁ (14.84). The variation in taste scores due to varieties was also significant. Variety V₂ had the highest taste score (18.35), followed by V₃ (17.59) and V₁ (17.22). The lowest taste score was recorded by variety V₅ (16.07). Cane regulation had a non-significant effect on taste during 2023, 2024, and in pooled data. In pooled data, cane C₃ exhibited the highest taste score (17.48), followed by C₂ (17.16) and control C₁ (16.93). The interaction between rootstocks, varieties, and cane regulation significantly influenced the taste evaluation. In pooled data, the treatment combination R₂V₂C₃ recorded the highest taste score (21.47), followed by R₂V₂C₂ (20.54) and R₂V₃C₃ (20.04).

Impact of Rootstock Selection and Cane Regulation on the Organoleptic Characteristics of Wine

Table 1: Organoleptic evaluation for appearance and aroma.

Treatment	Appearance			Aroma		
	2023	2024	Pooled	2023	2024	Pooled
Root Stocks (R)						
R ₁	13.51	12.38	12.94	14.10	12.74	13.42
R ₂	18.59	17.22	17.90	19.19	17.13	18.16
S.E. M±	0.02	0.04	0.01	0.04	0.04	0.02
CD AT 5%	0.14	0.27	0.06	0.25	0.25	0.10
Varieties (V)						
V ₁	16.41	14.69	15.55	17.11	14.99	16.05
V ₂	18.06	16.89	17.47	18.13	16.48	17.31
V ₃	17.12	15.80	16.46	17.44	15.82	16.63
V ₄	15.29	14.16	14.72	16.22	14.48	15.35
V ₅	13.36	12.47	12.91	14.32	12.92	13.62
S.E. M±	0.44	0.30	0.29	0.22	0.23	0.15
CD AT 5%	1.32	0.89	0.86	0.65	0.70	0.46
Canes (C)						
C ₁	15.71	14.47	15.09	16.23	14.66	15.44
C ₂	16.00	14.80	15.40	16.66	14.93	15.79
C ₃	16.43	15.13	15.78	17.04	15.22	16.13
S.E. M±	0.26	0.24	0.19	0.29	0.28	0.21
CD AT 5%	0.73NS	0.69NS	0.55	0.82NS	0.8NS	0.59NS
Root Stocks (R) x Canes (C) x Varieties (V)						
R ₁ V ₁ C ₁	13.84	11.89	12.86	14.51	12.66	13.59
R ₁ V ₁ C ₂	13.96	12.52	13.24	14.70	12.68	13.69
R ₁ V ₁ C ₃	14.10	13.02	13.56	14.80	12.90	13.85
R ₁ V ₂ C ₁	14.96	13.89	14.43	15.03	13.97	14.50
R ₁ V ₂ C ₂	15.06	14.52	14.79	15.59	14.29	14.94
R ₁ V ₂ C ₃	15.55	14.62	15.09	15.79	14.45	15.12
R ₁ V ₃ C ₁	14.46	13.38	13.92	14.82	13.60	14.21
R ₁ V ₃ C ₂	14.59	13.63	14.11	14.91	13.69	14.30
R ₁ V ₃ C ₃	14.86	13.82	14.34	14.93	13.78	14.36
R ₁ V ₄ C ₁	12.67	11.59	12.13	13.50	11.98	12.74
R ₁ V ₄ C ₂	12.89	11.64	12.27	13.58	12.57	13.07
R ₁ V ₄ C ₃	13.59	11.81	12.70	13.89	12.59	13.24
R ₁ V ₅ C ₁	10.37	9.11	9.74	10.38	10.10	10.24
R ₁ V ₅ C ₂	10.76	9.55	10.16	11.76	10.72	11.24
R ₁ V ₅ C ₃	10.93	10.72	10.82	13.32	11.16	12.24
R ₂ V ₁ C ₁	18.33	16.71	17.52	19.27	17.11	18.19
R ₂ V ₁ C ₂	18.91	16.97	17.94	19.61	17.27	18.44
R ₂ V ₁ C ₃	19.29	17.01	18.15	19.77	17.33	18.55
R ₂ V ₂ C ₁	20.18	19.22	19.70	20.55	18.44	19.50
R ₂ V ₂ C ₂	20.94	19.36	20.15	20.70	18.76	19.73
R ₂ V ₂ C ₃	21.66	19.70	20.68	21.13	18.97	20.05
R ₂ V ₃ C ₁	19.44	17.44	18.44	19.91	17.75	18.83
R ₂ V ₃ C ₂	19.52	17.94	18.73	19.95	17.93	18.94
R ₂ V ₃ C ₃	19.87	18.59	19.23	20.12	18.18	19.15
R ₂ V ₄ C ₁	17.20	16.55	16.87	18.44	16.41	17.43
R ₂ V ₄ C ₂	17.22	16.66	16.94	18.70	16.61	17.65
R ₂ V ₄ C ₃	18.17	16.68	17.43	19.18	16.70	17.94
R ₂ V ₅ C ₁	15.65	14.93	15.29	15.88	14.55	15.22
R ₂ V ₅ C ₂	16.19	15.23	15.71	17.09	14.79	15.94
R ₂ V ₅ C ₃	16.25	15.29	15.77	17.47	16.18	16.83
S.E. M±	0.81	0.76	0.61	0.91	0.88	0.66
CD AT 5%	2.31	2.17	1.74	2.60	2.52	1.88

Table 2: Organoleptic evaluation for taste and colour.

Treatment	Taste			Colour		
	2023	2024	Pooled	2023	2024	Pooled
Root Stocks (R)						
R ₁	15.49	14.19	14.84	16.30	15.11	15.70
R ₂	20.52	18.56	19.54	21.16	19.64	20.40
S.E. M±	0.05	0.04	0.02	0.05	0.04	0.04
CD AT 5%	0.30	0.26	0.14	0.32	0.25	0.24
Varieties (V)						
V ₁	17.88	16.55	17.22	18.57	17.23	17.90
V ₂	19.20	17.50	18.35	20.10	19.18	19.64
V ₃	18.31	16.88	17.59	19.48	17.69	18.59
V ₄	17.51	15.92	16.72	18.01	16.78	17.39
V ₅	17.11	15.04	16.07	17.48	16.02	16.75
S.E. M±	0.29	0.30	0.19	0.35	0.39	0.27
CD AT 5%	0.88	0.91	0.58	1.04	1.17	0.80
Canes (C)						
C ₁	17.75	16.11	16.93	18.49	17.05	17.77
C ₂	17.99	16.32	17.16	18.76	17.32	18.04
C ₃	18.27	16.70	17.48	18.94	17.76	18.35
S.E. M±	0.28	0.27	0.20	0.30	0.25	0.21
CD AT 5%	0.8NS	0.78NS	0.57NS	0.84NS	0.72NS	0.6NS
Root Stocks (R) x Canes (C) x Varieties (V)						
R ₁ V ₁ C ₁	15.10	14.30	14.70	16.00	14.70	15.35
R ₁ V ₁ C ₂	15.29	14.33	14.81	16.20	14.76	15.48
R ₁ V ₁ C ₃	15.56	14.38	14.97	16.67	14.76	15.72
R ₁ V ₂ C ₁	16.28	14.68	15.48	17.14	16.64	16.89
R ₁ V ₂ C ₂	16.49	14.99	15.74	17.55	16.99	17.27
R ₁ V ₂ C ₃	17.15	16.20	16.68	17.86	17.68	17.77
R ₁ V ₃ C ₁	15.62	14.38	15.00	16.77	14.87	15.82
R ₁ V ₃ C ₂	16.10	14.41	15.26	16.78	15.40	16.09
R ₁ V ₃ C ₃	16.18	14.67	15.42	16.90	15.77	16.33
R ₁ V ₄ C ₁	14.81	13.86	14.34	15.78	14.42	15.10
R ₁ V ₄ C ₂	14.97	13.92	14.45	15.84	14.54	15.19
R ₁ V ₄ C ₃	15.07	14.10	14.58	15.87	14.67	15.27
R ₁ V ₅ C ₁	14.38	12.51	13.45	14.47	13.03	13.75
R ₁ V ₅ C ₂	14.48	13.00	13.74	15.12	14.07	14.60
R ₁ V ₅ C ₃	14.80	13.19	14.00	15.51	14.34	14.93
R ₂ V ₁ C ₁	20.30	18.32	19.31	20.48	19.66	20.07
R ₂ V ₁ C ₂	20.50	18.93	19.72	20.98	19.71	20.34
R ₂ V ₁ C ₃	20.55	19.06	19.80	21.10	19.77	20.44
R ₂ V ₂ C ₁	21.06	19.33	20.20	22.29	20.65	21.47
R ₂ V ₂ C ₂	21.64	19.43	20.54	22.85	20.99	21.92
R ₂ V ₂ C ₃	22.56	20.38	21.47	22.95	22.09	22.52
R ₂ V ₃ C ₁	20.56	19.24	19.90	22.09	19.81	20.95
R ₂ V ₃ C ₂	20.64	19.24	19.94	22.12	19.82	20.97
R ₂ V ₃ C ₃	20.76	19.32	20.04	22.23	20.46	21.35
R ₂ V ₄ C ₁	19.95	17.68	18.81	20.13	18.85	19.49
R ₂ V ₄ C ₂	20.09	17.82	18.96	20.17	18.96	19.56
R ₂ V ₄ C ₃	20.20	18.12	19.16	20.28	19.22	19.75
R ₂ V ₅ C ₁	19.45	16.79	18.12	19.72	17.88	18.80
R ₂ V ₅ C ₂	19.72	17.16	18.44	19.99	17.95	18.97
R ₂ V ₅ C ₃	19.82	17.58	18.70	20.06	18.84	19.45
S.E. M±	0.89	0.86	0.63	0.93	0.79	0.66
CD AT 5%	2.54	2.47	1.81	2.67	2.27	1.88

Impact of Rootstock Selection and Cane Regulation on the Organoleptic Characteristics of Wine

Table 3: Overall organoleptic evaluation

Treatment	Overall organoleptic evaluation		
	2023	2024	Pooled
Root Stocks (R)			
R ₁	15.77	14.11	14.94
R ₂	20.41	18.87	19.64
S.E. M±	0.04	0.02	0.01
CD AT 5%	0.22	0.13	0.06
Varieties (V)			
V ₁	18.11	16.36	17.23
V ₂	19.58	17.60	18.59
V ₃	18.50	16.99	17.75
V ₄	17.47	15.99	16.73
V ₅	16.79	15.51	16.15
S.E. M±	0.28	0.27	0.22
CD AT 5%	0.84	0.81	0.65
Canes (C)			
C ₁	17.86	16.33	17.10
C ₂	18.06	16.50	17.28
C ₃	18.35	16.64	17.50
S.E. M±	0.25	0.28	0.21
CD AT 5%	0.71NS	0.81NS	0.6NS
Root Stocks (R) x Canes (C) x Varieties (V)			
R ₁ V ₁ C ₁	15.60	13.88	14.74
R ₁ V ₁ C ₂	15.80	13.91	14.85
R ₁ V ₁ C ₃	16.03	14.16	15.10
R ₁ V ₂ C ₁	16.80	15.21	16.00
R ₁ V ₂ C ₂	16.86	15.35	16.10
R ₁ V ₂ C ₃	17.27	15.40	16.34
R ₁ V ₃ C ₁	16.08	14.62	15.35
R ₁ V ₃ C ₂	16.13	14.68	15.41
R ₁ V ₃ C ₃	16.57	14.73	15.65
R ₁ V ₄ C ₁	14.88	13.45	14.16
R ₁ V ₄ C ₂	15.32	13.68	14.50
R ₁ V ₄ C ₃	15.39	13.75	14.57
R ₁ V ₅ C ₁	14.40	12.60	13.50
R ₁ V ₅ C ₂	14.73	13.08	13.91
R ₁ V ₅ C ₃	14.75	13.09	13.92
R ₂ V ₁ C ₁	20.23	18.65	19.44
R ₂ V ₁ C ₂	20.46	18.70	19.58
R ₂ V ₁ C ₃	20.53	18.86	19.69
R ₂ V ₂ C ₁	22.02	19.53	20.78
R ₂ V ₂ C ₂	22.18	19.76	20.97
R ₂ V ₂ C ₃	22.35	20.34	21.34
R ₂ V ₃ C ₁	20.54	19.14	19.84
R ₂ V ₃ C ₂	20.58	19.35	19.96
R ₂ V ₃ C ₃	21.12	19.42	20.27
R ₂ V ₄ C ₁	19.53	18.31	18.92
R ₂ V ₄ C ₂	19.74	18.35	19.04
R ₂ V ₄ C ₃	19.97	18.38	19.18
R ₂ V ₅ C ₁	18.57	17.91	18.24
R ₂ V ₅ C ₂	18.78	18.10	18.44
R ₂ V ₅ C ₃	19.50	18.30	18.90
S.E. M±	0.79	0.89	0.66
CD AT 5%	2.25	2.55	1.89

Organoleptic evaluation of colour

The colour evaluation in 2023, 2024, and pooled data revealed significant differences due to rootstocks (Table 2). In pooled data, rootstock R₂ recorded significantly higher colour scores (20.40) compared to rootstock R₁ (15.70). The variation in colour scores due to varieties was also significant. Variety V₂ had the highest colour score (19.64), followed by V₃ (18.59) and V₁ (17.90). The lowest colour score was recorded by variety V₅ (16.75). Cane regulation had a non-significant effect on colour during 2023, 2024, and in pooled data. In pooled data, cane C₃ exhibited the highest colour score (18.35), followed by C₂ (18.04) and control C₁ (17.77). The interaction between rootstocks, varieties, and cane regulation significantly influenced the colour evaluation. In pooled data, the treatment combination R₂V₂C₃ recorded the highest colour score (22.52), followed by R₂V₂C₂ (21.92) and R₂V₃C₃ (21.35).

Overall organoleptic evaluation

The overall organoleptic evaluation in 2023, 2024, and pooled data revealed significant differences influenced by rootstocks (Table 3). In pooled data, rootstock R₂ had a higher overall evaluation score (19.64) compared to R₁ (14.94), showing a notable difference in the quality of the treatments. For varieties, V₂ scored the highest in pooled data (18.59), followed by V₃ (17.75) and V₁ (17.23), with V₅ showing the lowest score (16.15). The effect of cane regulation on the overall evaluation was non-significant across all years and pooled data, as canes C₁, C₂, and C₃ displayed similar scores. In pooled data, C₃ had the highest score (17.50), followed by C₂ (17.28) and C₁ (17.10). The interaction between rootstocks, varieties, and cane regulation significantly affected the overall evaluation. In pooled data, the highest scores were observed in the combinations of R₂V₂C₃ (21.34), R₂V₂C₂ (20.97), and R₂V₂C₁ (20.78), with the lowest scores coming from combinations involving R₁, such as R₁V₃C₁ (13.50) and R₁V₄C₁ (14.16).

The pooled data demonstrated significant differences in organoleptic evaluations across rootstocks, varieties, and their interactions, showcasing their influence on grape quality parameters such as appearance, aroma, taste, and colour. Rootstock R₂ consistently outperformed R₁ in all parameters, ascribing its superior performance to enhanced nutrient uptake and better compatibility with scion varieties, corroborating findings by Tarricone *et al* (2023) who reported improved berry appearance and sensory quality in grapes grafted onto vigorous rootstocks. Similarly, the significant variation among

varieties highlights the influence of genetic factors, with variety V₂ excelling in appearance, aroma, taste, and colour. These results align with the studies of Singh *et al* (2022), where *Vitis vinifera* cultivars exhibited varietal differences in sensory traits due to variations in biochemical composition. The non-significant effects of cane regulation on aroma, taste, colour, and overall evaluation suggest that cane numbers alone may not have a direct influence on organoleptic properties. However, the superior scores observed in treatments involving C₃ could be attributed to its optimized canopy structure, leading to better light interception and photosynthetic efficiency, as discussed by Mian *et al* (2024). The interaction effects revealed that the highest scores in all organoleptic parameters were recorded for the R₂V₂C₃ combination, emphasizing the synergistic impact of selecting suitable rootstock-variety combinations alongside appropriate cane regulation. This finding aligns with reports by Blank *et al* (2022), who demonstrated that rootstock-scion interactions coupled with optimal cultural practices could significantly enhance grape quality.

CONCLUSION

The study highlighted the critical role of rootstocks and varietal characteristics in enhancing the organoleptic qualities of grapes, with R₂ and V₂ emerging as the most promising treatments. Additionally, while cane regulation had a limited direct impact, its interaction with rootstock and variety contributed significantly to the overall quality. These findings underline the importance of adopting an integrated approach combining rootstock selection, variety choice, and cultural practices to optimize grape production and sensory quality in the northern dry zone of Karnataka.

REFERENCES

- Blank M, Samer S and Stoll M (2022). Grapevine rootstock genotypes influences berry and wine phenolic composition (*Vitis vinifera* L. cv. Pinot Noir). *Oeno One* **56**(1): 133-144.
- Kaur J, Kaur G, Kaur K and Arora N K (2021). Yield and morphological characters as affected by chlormequat chloride application in grape cultivar Punjab MACS Purple. *JKrishi Vigyan* **10**(1): 314-320.
- Kiran Sagar D C, Athani S I, Gopali J B, Patil D R, Manukumar H R and Shivakumar K M (2019). Effect of gibberellic acid on berry quality parameters of different grape varieties under

Impact of Rootstock Selection and Cane Regulation on the Organoleptic Characteristics of Wine

- northern dry zone of Karnataka. *Int J Chem Stud* **7**(1): 759-762.
- Mian G, Colautti A, Belfiore N, Marcuzzo P, Tomasi D, Bell L. and Celotti E (2024). Fertigation affects photosynthesis, modulation of secondary metabolism and sensory profiles of *Vitis vinifera* cv. "Schioppettino" withered grapes and wines. *Sci Hort* **328**: 112954.
- Singh S, Bal J S and Singh J (2025). Performance of stionic combination on vegetative growth and flowering of exotic mandarin germplasm under sub-tropical plains of Punjab. *J Krishi Vigyan* **13**(1): 130-137.
- Singh B, Panesar P S and Nanda V (2022). Grape cultivar and drying method affect sensory characteristics. *Food Chem* **371**: 131162.
- Tarricone L, Amendolagine A M, Gennaro D, Masi G and Gentile G (2014). Rootstock effects on productive characteristics of "Princess" seedless table grapes in Apulia region (southern Italy). *International Symposium on Grapevine Roots* 1136 pp. 51-56.
- Joshi V, Kumar V, Debnath M, Pattanashetti S K, Variath M T and Khadakabhavi S (2015). Multivariate analysis of colored and white grape grown under semi-arid tropical conditions of Peninsular India. *Int J Agri Crop Sci* **8**(3): 350-365.

Received on 10/4/2025 Accepted on 12/5/2025