

Effect of Foliar Application of Micronutrients and Plant Growth Regulators on Quality of Guava (*Psidium Guajava* L.) Cv. Lalit

K Jawahar Srinith , CT Sathappan and Sharvesh S

Department of Horticulture, Faculty of Agriculture,
Annamalai University, Annamalai Nagar, 608002, Tamil Nadu, India

ABSTRACT

Lalit guava known for its distinct sweetness and vibrant red color is a favored variety among growers for its exceptional flavor and nutritional qualities. A field experiment was carried out during the year 2020–2021 on seven-year-old red guava trees at a farmer's orchard located in Semmangalam, Anathur, in the Villupuram district. The study aimed to identify the most suitable treatment for improving fruit quality in guava. The trial followed a Randomized Block Design with ten treatment combinations of T₁ (Control), T₂ (Boric acid @ 0.4% + GA₃ @ 150 ppm), T₃ (Chelated Zinc @ 0.5% + GA₃ @ 150 ppm), T₄ (Boric acid @ 0.4% + Chelated Zinc @ 0.5% + GA₃ @ 150 ppm), T₅ (Boric acid @ 0.4% + NAA @ 100 ppm), T₆ (Chelated Zinc @ 0.5% + NAA @ 100 ppm), T₇ (Boric acid @ 0.4% + Chelated Zinc @ 0.5% + NAA @ 100 ppm), T₈ (Boric acid @ 0.4% + SA @ 100 ppm) and T₉ (Chelated Zinc @ 0.5% + SA @ 100 ppm), T₁₀ (Boric acid @ 0.4% + Chelated Zinc @ 0.5% + SA @ 100 ppm). Each replicated three times. For optimal quality parameters such as total soluble solids, total sugars, reducing and non-reducing sugars, TSS/acid and minimal acidity, T₄ (Boric acid @ 0.4% + Chelated zinc @ 0.5% + GA₃ @ 150 ppm) exhibited superior results compared to the control. In contrast, T₇ (Boric acid @ 0.4% + Chelated Zinc @ 0.5% + NAA @ 100 ppm) was found to be most effective in enhancing the ascorbic acid and pectin content when compared to the untreated control.

Keywords: Fruit quality, Guava, Pectin content, Total sugar.

INTRODUCTION

Guava (*Psidium guajava* L.), a member of the Myrtaceae family, is extensively cultivated across tropical and subtropical regions. In India, it holds significant economic importance and is often dubbed the "Apple of the Tropics." This fruit is highly valued for its abundance of pectin and Vitamin C, and it also provides moderate levels of Vitamin A, calcium, phosphorus, pantothenic acid, riboflavin, thiamine, and niacin. Nutritionally, guava contains about 82.5% moisture, 2.45% acidity, 4.45% reducing sugars, 5.25% non-reducing sugars, 9.73 °Brix total soluble solids (TSS), and approximately 260 mg of Vitamin C per 100 grams of fresh fruit. It is especially suited for jelly production, and its juice is often used in ice cream formulations (Shahroon *et al.*, 2018). In addition to its nutritional profile, guava exhibits strong antioxidant activity due to its high Vitamin C levels and the presence of bioactive compounds such as carotenoids, isoflavonoids, and polyphenols. Although foliar sprays of micronutrients and plant growth regulators have been shown to improve fruit quality individually, there

is limited research on their synergistic effects in guava cultivation.

MATERIALS AND METHODS

The research entitled "Effect of Foliar Application of Micronutrients and Plant Growth Regulators on Quality of Guava (*Psidium guajava* L.) cv. Lalit" was carried out during the year 2020–2021 using seven-year-old red-fleshed guava trees. The study site was a farmer's field located in Semmangalam, Anathur, in the Villupuram district. A Randomized Block Design (RBD) was employed, comprising 10 treatment combinations with three replications each, involving a total of 300 guava plants. The trees were planted at a spacing of 9 × 9 meters to ensure adequate growth and canopy development. The foliar treatments of T₁ (Control), T₂ (Boric acid @ 0.4% + GA₃ @ 150 ppm), T₃ (Chelated Zinc @ 0.5% + GA₃ @ 150 ppm), T₄ (Boric acid @ 0.4% + Chelated Zinc @ 0.5% + GA₃ @ 150 ppm), T₅ (Boric acid @ 0.4% + NAA @ 100 ppm), T₆ (Chelated Zinc @ 0.5% + NAA @ 100 ppm), T₇ (Boric acid @ 0.4% + Chelated Zinc @ 0.5% +

Table 1. Effect of foliar application of micronutrients and plant growth regulators on total sugars, reducing sugar and non-reducing sugar of guava cv. Lalit (*Psidium guajava*).

Treatment	Details	TSS (°Brix)	Total sugars (%)	Reducing sugar (%)	Non- reducing sugar (%)
T ₁	Control	7.89	8.03	4.19	3.84
T ₂	Boric acid @ 0.4 % + GA ₃ @ 150 ppm	9.57	9.17	4.81	4.36
T ₃	Chelated zinc @ 0.5 % + GA ₃ @ 150 ppm	8.53	8.38	4.39	3.99
T ₄	Boric acid @ 0.4 % + Chelated zinc @ 0.5 % + GA ₃ @ 150 ppm	12.23	10.65	5.63	5.02
T ₅	Boric acid @ 0.4 % + NAA @ 100 ppm	11.59	10.31	5.43	4.88
T ₆	Chelated zinc @ 0.5 % + NAA @ 100 ppm	9.17	8.74	4.59	4.15
T ₇	Boric acid @ 0.4 % + Chelated zinc @ 0.5 % + NAA @ 100 ppm	10.95	9.96	5.23	4.73
T ₈	Boric acid @ 0.4 % + SA @ 100 ppm	10.21	9.52	5.01	4.51
T ₉	Chelated zinc @ 0.5 % + SA @ 100 ppm	9.37	9.09	4.79	4.30
T ₁₀	Boric acid @ 0.4 % + Chelated zinc @ 0.5 % + SA @ 100 ppm	10.85	9.87	5.21	4.66
S. ED		0.29	0.27	0.14	0.13
CD (p=0.05)		0.61	0.58	0.31	0.27

NAA @ 100 ppm), T₈ (Boric acid @ 0.4% + SA @ 100 ppm) and T₉ (Chelated Zinc @ 0.5% + SA @ 100 ppm), T₁₀ (Boric acid @ 0.4% + Chelated Zinc @ 0.5% + SA @ 100 ppm). consisting of various micronutrients and plant growth regulators, were administered at three stages: immediately after pruning, 30 days post-pruning, and 60 days after pruning. Data obtained from the experiment were analyzed statistically in accordance with the methodology proposed by Panse and Sukhatme (1961).

RESULTS AND DISCUSSION

The data (Table 1) indicated significant variations among treatments regarding parameters such as total soluble solids, total sugars, reducing and non-reducing sugars. Significantly higher Total Soluble Solids are observed with the treatment (T₄) with 12.23 °B over control (T₁) 7.89°B. The observed increase in Total Soluble Solids (TSS) may be attributed to the application of GA₃, which is known to enhance the accumulation of assimilates in developing fruits. GA₃ likely facilitates the mobilization of carbohydrates, promoting the conversion of complex polysaccharides into simpler sugars. Additionally, the role of boron in improving TSS could be linked to its involvement in transmembrane sugar transport. Boron is essential for several physiological and biochemical processes, including chlorophyll synthesis, enzymatic activities, respiration, and photosynthesis. Since sugars are the primary product of photosynthesis, an increase in photosynthetic activity due to boron application may lead to higher sugar synthesis, thereby contributing to greater TSS content in fruit juice (Ram and Bose, 2000). These results were in conformity with the results

reported by Kale *et al* (1999) in Ber. The increase in TSS has also been recorded by Yadav *et al* (2018), Lal and Das (2017) and Singh *et al* (2020) in guava.

The perusal of data on total sugars showed that the treatment T₄ (Boric acid @ 0.4 % + Chelated zinc @ 0.5 % + GA₃ @ 150 ppm) recorded maximum total sugars of 10.65 % over control (T₁) 8.03 %. This might be due to the fact that GA₃ increases the activity of the enzymes such as amylases, which hydrolyses the complex polysaccharides in to simple sugars.

The data on reducing sugars showed that the treatment T₄ (Boric acid @ 0.4 % + Chelated zinc @ 0.5 % + GA₃@ 150 ppm) recorded that maximum reducing sugar of 5.63 % over control (T₁) 4.19 %. This may be due to the fact that boron promotes cell division and development in the growth regions of the plant near the tips of shoots and roots.

Statistically significant variation was recorded for Non-reducing sugar with the application of Boric acid @ 0.4 % + Chelated zinc @ 0.5 % + GA₃ @ 150 ppm. The data on Non-reducing sugar showed that, the treatment T₄ recorded that maximum non-reducing sugar of 5.02 % over control (T₁) 3.84 %. The increase in sugar content could be attributed to the higher concentration of GA₃, which enhances the hydrolysis of starch into simpler sugars. The elevated levels of total sugars, including both reducing and non-reducing sugars, may result from the efficient translocation of photosynthates to the developing fruits, regulated by the action of GA₃. This hormone likely facilitates the mobilization of carbohydrates, contributing to improved sugar accumulation in the fruit. These findings are supported by the results obtained by

Effect of Foliar Application of Micronutrients and Plant Growth Regulators

Table 2. Effect of foliar application of micronutrients and plant growth regulators on acidity, Total Soluble Solids and TSS/acid ratio of guava cv. Lalit (*Psidium guajava*L.)

Treatment	Detail	Acidity (%)	TSS/Acid	Ascorbic acid (mg/100 g)	Pectin (%)
T ₁	Control	0.48	16.60	159.85	0.49
T ₂	Boric acid @ 0.4 % + GA ₃ @ 150 ppm	0.39	25.26	177.58	1.08
T ₃	Chelated zinc @ 0.5 % + GA ₃ @ 150 ppm	0.45	19.13	166.05	0.70
T ₄	Boric acid @ 0.4 % + Chelated zinc @ 0.5 % + GA ₃ @ 150 ppm	0.24	50.96	188.99	1.45
T ₅	Boric acid @ 0.4 % + NAA @ 100 ppm	0.27	42.93	194.15	1.66
T ₆	Chelated zinc @ 0.5 % + NAA @ 100 ppm	0.42	22.02	171.21	0.86
T ₇	Boric acid @ 0.4 % + Chelated zinc @ 0.5 % + NAA @ 100 ppm	0.31	35.32	200.35	1.87
T ₈	Boric acid @ 0.4 % + SA @ 100 ppm	0.36	28.36	182.74	1.24
T ₉	Chelated zinc @ 0.5 % + SA @ 100 ppm	0.40	24.13	176.37	1.03
T ₁₀	Boric acid@ 0.4 % + Chelated zinc @ 0.5 % + SA @ 100 ppm	0.32	33.91	187.90	1.41
S. ED		0.29	0.01	0.87	5.27
CD (p=0.05)		0.61	0.02	1.83	11.07

Agrawal and Dikshit (2010) in sapota and Garasiya *et al* (2013) in guava.

The data presented in Table 2 Statistically significant variation was recorded for acidity with the application of micronutrients and plant growth regulators. A closer observation of data on acidity showed that, the treatment T₄ (Boric acid @ 0.4 % + Chelated zinc @ 0.5 % + GA₃150 ppm) recorded as minimum acidity of 0.24 % over control (T₁) 0.45 %. The reduction in acidity content may be attributed to the influence of boron, which becomes neutralized through plant metabolic pathways and is also utilized as a substrate in respiratory processes. The decreased acidity observed in the fruits could also be linked to the advancement of ripening triggered by the application of micronutrients, during which the degradation of organic acids likely takes place. These results are in accordance with the findings of Singh *et al* (2017) and Dutta and Banik (2005) in guava.

The data on ascorbic acid showed that, the treatment T₇ (Boric acid @ 0.4 % + Chelated zinc @ 0.5 % + NAA @ 100 ppm) recorded as maximum ascorbic acid content (200.35 mg) over control T₁ (159.85 mg). An increased in vitamin C content was due to catalytic influence of NAA on bio synthesis of vitamin C.

or both. Similar trend was also observed by Brahmachari *et al* (1997), Dubey *et al* (2002) and

Garasiya *et al* (2013) in guava.

The data on pectin content showed that, the treatment T₇ (Boric acid @ 0.4 % + Chelated zinc @ 0.5 % + NAA @ 100 ppm) recorded that maximum pectin content of 0.24 % over control (T₁) 0.49 %. This effect may be attributed to the application of NAA, which facilitates the solubilization of pectic substances from the middle lamella, thereby contributing to an increase in pectin content. Kumar *et al.* (2010) also observed that the application of NAA in guava proved effective in increasing pectin content. Similar trend was also observed by Suman *et al* (2021) and Kumari *et al* (2021).

CONCLUSION

The study highlighted the significant influence of micronutrients and plant growth regulators on fruit quality parameters. Treatment T₄ (Boric acid @ 0.4% + Chelated zinc @ 0.5% + GA₃ @ 150 ppm) proved most effective in enhancing total soluble solids, total sugars, reducing sugars, and non-reducing sugars while reducing acidity. Meanwhile, T₇ (Boric acid @ 0.4% + Chelated zinc @ 0.5% + NAA @ 100 ppm) significantly increased ascorbic acid and pectin content. These findings suggested that the combined application of micronutrients and PGRs can effectively improve fruit quality, warranting further research for optimized formulations.

REFERENCE

- Agrawal S and Dikshit SN (2010). Studies on the effect of plant growth regulators on qualitative characters of sapota cv. Cricket Ball. *Indian J Hort* **67(2)**: 177-180.
- Brahmachari VS, Kumar N and Kumar R (1997). Effect of foliar feeding of calcium, Potassium and growth substances on yield and quality of guava (*Psidium guajava* L.). Haryana. *J Hort Sci* **26 (3-4)**: 169-173.
- Dubey AK, Singh DB, Barhe S, Singh A and alal M (2002). Deblossoming in summer season flowering in guava. *Indian Hort* **47(1)**: 35-36
- Dutta P and Banik AK (2005). Effect of foliar feeding of nutrients and plant growth regulators on physico-chemical quality of sardar guava grown in red and Lateritic tract of West Bengal. *Int Symposium on guava*. **735**: 407-411.
- Garasiya VR, Patel NM, Bhadauria HS and Wankhade VR (2013). Effect of plant growth regulators on quality of winter season guava (*Psidium guajava* L.) cv. Lucknow-49. *Asian J Hort* **8**: 347-9
- Kale VS, Kale SPB and Adpawar RM (1999). Effect of plant growth regulators on fruit yield quality of ber cv. *Umrana* (*Zizyphus mauritiana* Lamk.). *Annual Revi Pl Physiol* **13(1)**: 69-72.
- Kumar M, Kumar R and Singh RP (2010). Effect of micronutrients and plant growth regulators on fruiting of litchi. *Indian J Agri Sci* **5(2)**: 521-524.
- Kumari P, Prasad VM, Bahadur V and Sahoo R (2021). Effect of different micronutrient on plant growth, yield and fruit quality in Existing Guava (*Psidium guajava* L.) cv. Allahabad Safeda. *Pharma Innov* **10(9)**: 316-320.
- Lal N, Das RP (2017). Effect of plant growth regulators on yield and quality of guava (*Psidium guajava* L.) cv. Allahabad Safeda. *Int J Curr Microbiol Appl Sci*. **6(5)**: 857-863.
- Mandal G, Dhaliwal HS and Mahajan BVC (2012). Effect of pre-harvest application of NAA and potassium nitrate on storage quality of winter guava (*Psidium guajava*). *Indian J Agri Sci* **82(11)**.
- Panse VG, Sukhatme PV (1954). Statistical methods for agricultural workers. Indian Council of Agricultural Research (ICAR), New Delhi.
- Ram RA and, Bose TK (2000). Effect of foliar application of magnesium and micronutrients on growth, yield and fruit quality of mandarin orange (*Citrus reticulata* Blanco). *Indian J Hort* **57(3)**: 215-220.
- Shahroon Khan, Kumar A, Dahiya DS, Baloda S, Malik A (2018). Significance of nutrient application on growth, yield and quality of Guava: A review. *Int J Chem Stud* **6(3)**: 2936-2942
- Singh J and Mirza A (2017). Study on physiochemical properties of fruits as influenced by naphthalene acetic acid: a review. *Int J Eng Res and Technol* **6**: 941-945.
- Singh, Uma Kant and Choudhary Ashwini (2020). Effect of foliar spray of plant growth regulators and micronutrient on guava (*Psidium guajava*). *Int J Agriworld* **1(1)**: 8-13
- Suman A, Singh, A. K. Kanth N and Kumar A (2021). Effect of foliar feeding of micronutrients and plant growth regulators on flowering and physical parameters of guava (*Psidium guajava* L.) cv. Allahabad Safeda under agroecological conditions of North Bihar. *Pharma Innov J* **10(9)**: 483-86.
- Yadav A, Ram RB, Verma RS, Kumar V and Kewal Yadav R (2018). Effect of foliar application of micronutrients on Bio-chemical attributes of winter season guava (*Psidium guajava* L.) cv. Lalit. *J Pharmaco and phytochem* **7(2)**, pp.3196-3197.

Received on 11/4/2025 Accepted on 13/5/2025